

The 60th William Lowell Putnam Mathematical Competition
Saturday, December 4, 1999

A-1 Find polynomials $f(x)$, $g(x)$, and $h(x)$, if they exist, such that for all x ,

$$|f(x)| - |g(x)| + h(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0. \end{cases}$$

A-2 Let $p(x)$ be a polynomial that is nonnegative for all real x . Prove that for some k , there are polynomials $f_1(x), \dots, f_k(x)$ such that

$$p(x) = \sum_{j=1}^k (f_j(x))^2.$$

A-3 Consider the power series expansion

$$\frac{1}{1-2x-x^2} = \sum_{n=0}^{\infty} a_n x^n.$$

Prove that, for each integer $n \geq 0$, there is an integer m such that

$$a_n^2 + a_{n+1}^2 = a_m.$$

A-4 Sum the series

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{m^2 n}{3^m (n3^m + m3^n)}.$$

A-5 Prove that there is a constant C such that, if $p(x)$ is a polynomial of degree 1999, then

$$|p(0)| \leq C \int_{-1}^1 |p(x)| dx.$$

A-6 The sequence $(a_n)_{n \geq 1}$ is defined by $a_1 = 1$, $a_2 = 2$, $a_3 = 24$, and, for $n \geq 4$,

$$a_n = \frac{6a_{n-1}^2 a_{n-3} - 8a_{n-1} a_{n-2}^2}{a_{n-2} a_{n-3}}.$$

Show that, for all n , a_n is an integer multiple of n .

B-1 Right triangle ABC has right angle at C and $\angle BAC = \theta$; the point D is chosen on AB so that $|AC| = |AD| = 1$; the point E is chosen on BC so that $\angle CDE = \theta$. The perpendicular to BC at E meets AB at F . Evaluate $\lim_{\theta \rightarrow 0} |EF|$.

B-2 Let $P(x)$ be a polynomial of degree n such that $P(x) = Q(x)P''(x)$, where $Q(x)$ is a quadratic polynomial and $P''(x)$ is the second derivative of $P(x)$. Show that if $P(x)$ has at least two distinct roots then it must have n distinct roots.

B-3 Let $A = \{(x, y) : 0 \leq x, y < 1\}$. For $(x, y) \in A$, let

$$S(x, y) = \sum_{\frac{1}{2} \leq \frac{m}{n} \leq 2} x^m y^n,$$

where the sum ranges over all pairs (m, n) of positive integers satisfying the indicated inequalities. Evaluate

$$\lim_{(x,y) \rightarrow (1,1), (x,y) \in A} (1 - xy^2)(1 - x^2y)S(x, y).$$

B-4 Let f be a real function with a continuous third derivative such that $f(x)$, $f'(x)$, $f''(x)$, $f'''(x)$ are positive for all x . Suppose that $f'''(x) \leq f(x)$ for all x . Show that $f'(x) < 2f(x)$ for all x .

B-5 For an integer $n \geq 3$, let $\theta = 2\pi/n$. Evaluate the determinant of the $n \times n$ matrix $I + A$, where I is the $n \times n$ identity matrix and $A = (a_{jk})$ has entries $a_{jk} = \cos(j\theta + k\theta)$ for all j, k .

B-6 Let S be a finite set of integers, each greater than 1. Suppose that for each integer n there is some $s \in S$ such that $\gcd(s, n) = 1$ or $\gcd(s, n) = s$. Show that there exist $s, t \in S$ such that $\gcd(s, t)$ is prime.